



# Carbon and nitrogen mineralization and enzyme activities in soil aggregate-size classes: Effects of biochar, oyster shells, and polymers

Yasser Mahmoud Awad <sup>a, b, c</sup>, Sang Soo Lee <sup>e</sup>, Ki-Hyun Kim <sup>d</sup>, Yong Sik Ok <sup>b, \*</sup>, Yakov Kuzyakov <sup>f, g</sup>

<sup>a</sup> Department of Soil Science of Temperate Ecosystems, Department of Agricultural Soil Science, University of Göttingen, 37077 Göttingen, Germany

<sup>b</sup> Korea Biochar Research Center, O-Jeong Eco-Resilience Institute (OJERI) & Division of Environmental Science and Ecological Engineering, Korea University, Seoul 02841, Republic of Korea

<sup>c</sup> Faculty of Agriculture, Suez Canal University, Ismailia 41522, Egypt

<sup>d</sup> Department of Civil and Environmental Engineering, Hanyang University, Seoul 04763, Republic of Korea

<sup>e</sup> Division of Environmental Engineering, Yonsei University, Wonju, Republic of Korea

<sup>f</sup> Department of Agricultural Soil Science, University of Göttingen, 37077 Göttingen, Germany

<sup>g</sup> Institute of Environmental Sciences, Kazan Federal University, 420049 Kazan, Russia

## HIGHLIGHTS

- Oyster shell (OS), biochar (BC), and biopolymer (BP) changed N-cycling in soil.
- Addition of OS to BC and to BP increased NO<sub>3</sub> in soil.
- OS alone or with BC or BP increased soil chitinase and aminopeptidase activities.
- BC/BP and OS contributes to soil fertility improvements and C sequestration.

## ARTICLE INFO

### Article history:

Received 10 November 2017

Received in revised form

4 January 2018

Accepted 10 January 2018

Available online 12 January 2018

Handling Editor: Patryk Oleszczuk

### Keywords:

Black carbon

Polymers

Soil aggregation

Nutrient dynamics

Enzyme activities

## ABSTRACT

Biochar (BC) and polymers are cost-effective additives for soil quality improvement and long-term sustainability. The additional use of the oyster shells (OS) powder in BC- or polymer-treated soils is recommended as a nutrient source, to enhance aggregation and to increase enzyme activities. The effects of soil treatments (i.e., BC (5 Mg ha<sup>-1</sup>) and polymers (biopolymer at 0.4 Mg ha<sup>-1</sup> or polyacrylamide at 0.4 Mg ha<sup>-1</sup>) with or without the OS (1%) on the short-term changes were evaluated based on a 30-day incubation experiment with respect to several variables (e.g., CO<sub>2</sub> release, NH<sub>4</sub><sup>+</sup> and NO<sub>3</sub><sup>-</sup> concentrations, aggregate-size classes, and enzyme activities in an agricultural Luvisol). The BC and BP with the addition of OS increased the portion of microaggregates (<0.25 mm) relative to the control soil without any additions, while PAM alone increased the portion of large macroaggregates (1–2 mm). Concentrations of NO<sub>3</sub><sup>-</sup> also increased in soils treated with OS, OS + BC, and OS + BP as result of the increased chitinase and leucine aminopeptidase activities. The BC and BP when treated with the additional OS had significant short-term impacts on N mineralization without affecting C mineralization in soil. Consequently, the combination of BC or BP with OS was seen to accelerate N turnover without affecting C turnover (and related C losses) from soil. As such, the addition of these additives contributed considerably to the improvement of soil fertility and C sequestration.

© 2018 Elsevier Ltd. All rights reserved.

## 1. Introduction

The deficiency of C and N limits the quality and productivity of

many soils (Lee et al., 2009; Katsalirou et al., 2010). Mineralization of C and N through decomposition of soil organic matter (SOM) exerts dominant controls on nutrient bioavailability (Weintraub and Schimel, 2003; Rothlisberger-Lewis et al., 2016). Furthermore, soil aggregates alter SOM decomposition as microaggregates may occlude SOM, and render it inaccessible to microbial decomposition (Awad et al., 2013; Kuzyakov et al., 2014). The decomposition of

\* Corresponding author.

E-mail address: [yongsikok@korea.ac.kr](mailto:yongsikok@korea.ac.kr) (Y.S. Ok).